arXiv:1204.5014v1 [math.MG] 23 Apr 2012

# Malfatti's problem on the hyperbolic plane*

Ákos G.Horváth

April 20, 2012

**Abstract**

More than two centuries ago Malfatti (see [10]) raised and solved the following problem called by Malfatti's construction problem: In a triangle describe three small circles, each of them touching the other two, and also two sides of the triangle. Interesting fact that nobody investigated this problem on the hyperbolic plane, while the case of the sphere were solved at once. We now compensate for this shortage and solve the following exercise: *Determine three cycles of the hyperbolic plane, each of them touching the two others, and also two of three given cycles of the hyperbolic plane.* We also give a proof of the hyperbolic version of Malfatti's marble problem.

**MSC(2000):** 51M10, 51M15

**Keywords:** cycle, hyperbolic plane, inversion, Malfatti's construction problem, Malfatti's marble problem,

## 1 Malfatti's construction problem

### 1.1 Once more again on the history of the problem

Malfatti (see [10]) raised and solved the following problem called by Malfatti's construction problem: *In a triangle describe three small circles, each of them touching the other two, and also two sides of the triangle.* Malfatti determined the radiuses of the required circles and thus he got an analytic-geometric solution. This formula on the radii was only the beginning of a discussion motivated by the original problem. We would like to mention here some nice further investigations.

The first incredible moment was Steiner's construction. He gave an elegant method (without proof) to find the Malfatti's circles and extend the problem and his construction to the case of circles (see in [14], [15]). Cayley in [3] referred to this problem as Steiner's extension of Malfatti's problem: *Determine three circles of the Euclidean plane, each of them touching the two others, and also two of three given circles.* We note that Cayley investigated and solved the

---

[*] Dedicated to the memory of my colleague, patron and friend István Reiman.

generalization of it in [3], he called it also by Steiner's extension of Malfatti's problem. His problem is: *Determine three sections of a surface of the second order, each of them touching the two others, and also two of three given sections of the surface of the second order.* In his paper showed that the Steiner's construction can be interpreted on a surface of second order and solved the corresponding problem of construction. Since the case of circles on the sphere is a generalization of the case of circles of the plane (as it can be seen easily by stereographic projection) Cayley indirectly proved also the Steiner's second construction. We also have to mention here the nice geometric proof of Hart for the Steiner's construction which published in [9]. (It can be found in various textbooks e.g. [6] and also on the web.)

The second moment which I would like to mention here two short papers written by Dr.Schellbach [12] and [13]. He solved the original construction problem and its "spherical variation", too. The elegant goniometric solution determines the touching points of the circles on the sides of the triangle (spherical triangle). These nice and surprising papers inspire Cayley to write a paper [4] in which he suggested a change of notation by which he did the Schellbach's formulas a little bit more elegant. As a consequence he supposed that the sides become indefinitely small and he had the case of plane triangle. The formulas led to some formulas of his third previous paper on this problem [5] and finally he determined the distances of the points of contact from the adjacent angles of the triangle by the side-lengthes of the triangle. He noted also that the equations are very similar in form to those given in the same paper for the determination of the radii of the inscribed Malfatti's circles. At this point seems to be that there are not possibility to give any new observations for this problem. However in this twenty-one's century Bottema (in [2]) observed a simple solution by inversion in the case when the determining figures are touching circles on the euclidean plane. The idea of the construction is that an inversion with respect to a circle which center is a common touching point of two given circle leads such a configuration in which the resulting circles can be got in a simple way. From this configuration can be calculated the radiuses of the resulting circles by the radiuses of the given circles.

Interesting fact that nobody consider the hyperbolic case of this – more than two centuries ago raised – problem, while the case of the sphere were solved at once. We now compensate for this shortage.

Additionally we give a proof for the hyperbolic case of the marble problem, which is the another problem corresponding to the original Malfatti's paper. Agree with Remark 2 in [1] we prove that: *On the hyperbolic plane the greedy arrangement has the largest total area among of non-overlapping circles in a triangle.*

## 1.2 The case of hyperbolic triangle

Note that the case of the hyperbolic triangle can be solved immediately (and independently for any model) by the method of Steiner and also the method of Schellbach, respectively. The exercise now is the following: *In a hyperbolic* *triangle describe three small circles, each of them touching the other two, and also two sides of the triangle.*

### 1.2.1 The Steiner's construction for hyperbolic triangle

To prove Malfatti's problem Steiner proposed the following construction [15]:

1. Draw three angle bisectors $OA$, $OB$ and $OC$. In the triangles $\triangle_{OAB}$, $\triangle_{OBC}$, $\triangle_{OCA}$ inscribe circles $c_C$, $c_A$, $c_B$, respectively.
2. For each pair of the circles consider the second (distinct to the corresponding angle bisector) internal tangents. The latter concur in a point $K$ and cross the sides in points $H, I$; $D, E$; and $F, G$; respectively.
3. The three quadrilaterals $KHCI$, $KGBH$, and $KEAG$ are inscriptible. Their incircles solve Malfatti's problem.

The proof of Hart [9] works on the hyperbolic plane, too. It is based on two elementary but absolute lemmas, implying that their corollaries are also true on the hyperbolic plane. These are:

**Lemma 1** *If the sum or difference of two tangents drawn from a point $P$ to two circles be equal to a common tangent of the circles, the point $P$ is on a common tangent.*

Hence it is evident that if the common tangents to each pair of three circles pass through the same point, one of these common tangents must be equal to the sum of other two, and that therefore the other common tangents will also pass through one point.

**Lemma 2** *If two circles cut off equal parts from a given right line and if tangents at the extreme points $A$, $B$ intersect at $P$, the circles will subtend equal angles at $P$, and also that if tangents be drawn from each point $A$ and $B$ to the other circle, they will be equal.*

The proof of the first lemma based on two absolute fact on the triangle inequality and on that the tangents drawing from a point to a circle are equals, respectively. The proof of the second one use trigonometry and can be interpreted in the hyperbolic plane, too. Hart himself observed that his proof works on the sphere, too. The Hart's proof of the construction (which we are citing word-by-word from the original paper with respect to its elegancy and clarity) is the following:

LET $L$, $M$, $N$, BE THE POINTS OF CONTACT OF THREE CIRCLES WHICH TOUCH ONE ANOTHER, AND EACH TOUCH TWO SIDES OF THE GIVEN TRIANGLE $\triangle_{ABC}$. DRAW $DE$, $FG$, $HI$, TOUCHING THESE CIRCLES AT $L, M, N$ AND MEETING ONE ANOTHER AT $K$. THEN SINCE $FH - HD = FO - DP = FM - DL = FK - DK$, $H$ IS THE POINT OF CONTACT OF THE CIRCLE INSCRIBED IN THE TRIANGLE $\triangle_{DKF}$; IN LIKE MANNER $E$ AND $G$ ARE THE POINTS OF CONTACT OF CIRCLES WHICH TOUCH $IK$, $KF$, AND $AC$; $IK$, $KD$,



Figure 1: Hart's proof of Steiner's construction.

AND $AB$, RESPECTIVELY; BUT $HN = HP = QL$, AND $NS = ER = EL$, THEREFORE $HS = EQ$, AND THEREFORE THE CIRCLES $HQ$ AND $ES$ SUBTEND EQUAL ANGLES AT $C$ (LEMMA 2). ALSO THREE COMMON TANGENTS $QL$, $SN$, AND $KF$ OF THE CIRCLES $HQ$, $ES$, $PNR$, PASS THROUGH $K$; THEREFORE $C$ MUST BE A POINT ON THE OTHER COMMON TANGENT TO $HQ$ AND $ES$ (COR. LEMMA 1). IN LIKE MANNER IT IS PROVED THAT THE BISECTORS OF $A$ AND $B$ ARE COMMON TANGENTS TO $ES$ AND $GT$; $HQ$ AND $GT$ RESPECTIVELY. WHENCE STEINER'S CONSTRUCTION IS EVIDENT FOR PLANE AND SPHERICAL TRIANGLES.

By inversion Hart extends his proof to all configurations of the plane or the sphere in which the sides of the given triangle are arcs of circles. We note that inversion could not change intersecting circles into non-intersecting one thus this method unable to prove Steiner's extension of Malfatti's problem in the case when the given circles are pairwise disjoints.

### 1.2.2 Schellbach's solution

The Schellbach's solution is a goniometric construction. Historically, this solution preceded the Hart's proof (see [12] and [13] ). To use this for hyperbolic case we change the Schellbach's equalities on spherical triangles onto the equalities of hyperbolic triangles. Practically we substitute imaginary values into the trigonometric function and change to the corresponding hyperbolic functions. Formally here we follow Cayley's simplified terminology. Let the sides of the triangle be $a, b, c$ and let $x, y, z$ be the distances of the points of contact from the adjacent angles of the triangle. Then writing

$$a + b + c = 2s,$$

$$a - \frac{1}{2}s = l, \quad b - \frac{1}{2}s = m, \quad c - \frac{1}{2}s = n$$

whence $l+m+n=\frac{1}{2}s$, and putting also

$$\frac{1}{2}s-x=\xi,\quad \frac{1}{2}s-y=\eta,\quad \frac{1}{2}s-z=\zeta,$$

then we have

$$\begin{aligned}
\frac{\cosh l\cosh\eta\cosh\zeta}{\cosh\frac{1}{2}s}+\frac{\sinh l\sinh\eta\sinh\zeta}{\sinh\frac{1}{2}s} &= 1\\
\frac{\cosh m\cosh\zeta\cosh\xi}{\cosh\frac{1}{2}s}+\frac{\sinh m\sinh\zeta\sinh\xi}{\sinh\frac{1}{2}s} &= 1\\
\frac{\cosh n\cosh\xi\cosh\eta}{\cosh\frac{1}{2}s}+\frac{\sinh n\sinh\xi\sinh\eta}{\sinh\frac{1}{2}s} &= 1,
\end{aligned}$$

from which equations the unknown quantities $\xi,\eta,\zeta$ are to be determined. To solve the equations, let the subsidiary angles $\lambda,\mu,\nu$ be determined by the conditions

$$\begin{aligned}
\frac{\cosh\lambda\cosh m\cosh n}{\cosh\frac{1}{2}s}-\frac{\sinh\lambda\sinh m\sinh n}{\sinh\frac{1}{2}s} &= 1\\
\frac{\cosh\mu\cosh n\cosh l}{\cosh\frac{1}{2}s}-\frac{\sinh\mu\sinh n\sinh l}{\sinh\frac{1}{2}s} &= 1\\
\frac{\cosh\nu\cosh l\cosh m}{\cosh\frac{1}{2}s}-\frac{\sinh\nu\sinh l\sinh m}{\sinh\frac{1}{2}s} &= 1,
\end{aligned}$$

then it may be shown that

$$\cosh(\eta+\zeta)=\frac{\cosh\left(\frac{s+\lambda-l}{2}\right)}{\cosh\left(\frac{\lambda+l}{2}\right)},\quad \cosh(\eta-\zeta)=\frac{\cosh\left(\frac{s-\lambda+l}{2}\right)}{\cosh\left(\frac{\lambda+l}{2}\right)},$$

$$\cosh(\zeta+\xi)=\frac{\cosh\left(\frac{s+\mu-m}{2}\right)}{\cosh\left(\frac{\mu+m}{2}\right)},\quad \cosh(\zeta-\xi)=\frac{\cosh\left(\frac{s-\mu+m}{2}\right)}{\cosh\left(\frac{\mu+m}{2}\right)},$$

$$\cosh(\xi+\eta)=\frac{\cosh\left(\frac{s+\nu-n}{2}\right)}{\cosh\left(\frac{\nu+n}{2}\right)},\quad \cosh(\xi-\eta)=\frac{\cosh\left(\frac{s-\nu+n}{2}\right)}{\cosh\left(\frac{\nu+n}{2}\right)}.$$

If we write

$$\begin{aligned}
\tanh\phi &= \tanh m\tanh n\coth\frac{1}{2}s\\
\tanh\chi &= \tanh n\tanh l\coth\frac{1}{2}s\\
\tanh\psi &= \tanh l\tanh m\coth\frac{1}{2}s,
\end{aligned}$$

then

$$\cosh(\lambda-\phi) = \frac{\cosh\frac{1}{2}s\cosh\phi}{\cosh m}\cosh n$$

$$\cosh(\mu-\chi) = \frac{\cosh\frac{1}{2}s\cosh\chi}{\cosh n}\cosh l$$
$$\cosh(\nu-\psi) = \frac{\cosh\frac{1}{2}s\cosh\psi}{\cosh l}\cosh m,$$

equations which give the values of $\lambda, \mu, \nu$ by the help of hyperbolic rectangular triangles. From these quantities $\xi, \eta, \zeta$ and thus $x, y, z$ can be constructed, too.

## 1.3 The case of the cycles

Our goal to investigate touching cycles of the hyperbolic plane. For simplicity in further we assume that **the domains of the given cycles are pairwise non-overlapping.** We formulate the exercise as follows:

*Determine three cycles of the hyperbolic plane, each of them touching the two others, and also two of three given cycles of the hyperbolic plane.*



Figure 2: The euclidean solution does not work on the hyperbolic plane.

Using Poincare's conformal disk model the given cycles represented by the arcs of certain circles. By Steiner's method we can construct the corresponding Malfatti's circles solving the Steiner's extension of Malfatti's problem. But may be possible that the corresponding arcs of circles does not intersect the model-circle meaning that there is no solution with respect to the hyperbolic plane. In fact, consider the situation of Fig.3. The edges of the regular triangle with respect to the basic circle of the Poincare's model represents arcs of hypercycles of the hyperbolic plane. On the other hand the uniquely determined external Malfatti's circles with respect to this regular triangle touche the corresponding edges in external points of the model implying that the corresponding cycles cannot be touching cycles to the given hypercycles.

The trouble of this example can be solved easily if we consider the whole cycle without an arc of a cycle. We will use the following concepts:

**Definition 1** *The* hypercycle *is the boundary of the locus of points of the plane with distances from a line is less or equal to a constant. (Specially it is a convex curve with two connected components.)* Two cycles are touching *if they have a common point with a common tangent line.*

We can state the following theorem on existence.

**Theorem 1** *For three given cycles of the hyperbolic plane has Malfatti's system of cycles.*

Remark that in this theorem on existence the domains of the corresponding cycles need not to be disjoint. On the other hand the tools of a concrete construction will require some further geometric conditions thus here also may assume that the domains of the given cycles are pairwise non-overlapping.
**Proof:** Consider the Poincare's disk model of the hyperbolic plane. In the case, when there is no hypercycle among the given cycles the statement is trivial. On the other hand by our definition a hypercycle is the union of two arcs of circles. Represent now a hypercycle with one of the whole circle containing one of the original arcs. The inner arc (which lies in the interior of the model) of the chosen circle is the inverse image (with respect to the model circle) of the outer arc of the other possible circle. Now Steiner's construction can be done on the embedding Euclidean plane and we get a Malfatti's system of circles for the representing circles. If we take the inverse of those points of the six circles which are external to the model, we get either whole hypercycle; paracycle; or a circle of the hyperbolic plane accordingly to the cases where the getting circle intersects; touched externally; or lies in the complement of the model circle.

So with this transformation from a Malfatti's system of circles of the embedding Euclidean plane, we get a Malfatti's system of cycles of the hyperbolic plane corresponding to the given cycles as we stated. □

Unfortunately the using condition for touching cycles cannot exclude the possibility of existences of further common points of the two cycles (contrary to the case of circles). Two possible points of intersection can lie on the other connected components of the hypercycle. (On the example of Fig.1 we can find this situation after the inversion of the external parts of the hypercycles.)

Thus with more rigorous definitions of touching we have new (typically harder) problems on existence. Two possibilities for the concepts of touching are:

- two cycles are touching if they have exactly one common (real) points on the hyperbolic plane (in that point they tangents are also the same), or
- two cycles are touching if they domains are touching externally.

In this paper we do not deal with these questions which can give the base of a foregoing paper. We consider here the most simple case of touching as we fixed it in our definition Definition 1. Our classical question is:

*How can we construct the Malfatti's cycles of a given system of cycles?*

The question is interesting if we cannot use model representations. Our purpose is to give such a construction. To this we have to build the hyperbolic analogue of some Euclidean concepts.

### 1.3.1 Hyperbolic power, inversion and centres of similitude on the hyperbolic plane

It is not clear that who investigated first the concept of inversion with respect to hyperbolic geometry. A synthetic approach can be found in [11] using reflections in Bachmann's metric plane. To our purpose more convenient an analytic building up in which also can define the concepts of centres of similitude and axe of similitude, respectively. We consider – as an analogy – the following spherical building up of these concepts.

It can be proved that if an arc of a great circle (line) passing through a fixed point $O$ cut a small circle in the points $A$,$B$,

$$\tan\frac{1}{2}OA\cdot\tan\frac{1}{2}OB$$

is constant. This product is called the *spherical power* of $O$ with respect to circle. It is positive or negative, according as $O$ is exterior or interior to the circle. If from any point $O$ outside a small circle two arcs be drawn to it, of which one, $OD$, is a tangent, and the other a secant, meeting it in the points $A,B$; then

$$\tan^2\frac{1}{2}OD=\tan\frac{1}{2}AO\cdot\tan\frac{1}{2}OB.$$

If we have two small circles on the sphere and consider the locus of $P$ for which the tangents to these circles are equals is a great circle called by the *radical circle (axe of power)* of them. The radical circles of three small circle taken in pairs are concurrent. The common point is the *power point* of the three small circle. This is the centre of the circle orthogonal to each of them.

For two small circles there are two *centres of similitude.* These are the points on the line connecting their centres which divide the segment joining the centers of the two circles externally and internally in the spherical ratio of the sine of the radii. Common tangents to the circles pass through the centres of similitude, viz., the direct common tangents through the external centre and the inverse common tangent through the internal centre. If through a centre of similitude we draw a secant cutting the circles, then the pairs of points $M,M'$;$N,N'$ of Fig.2 are said to be homothetic and $M,N'$; $M',N$ are inverse.

Then for the homothetic points $M,M'$

$$\tan\frac{SM}{2}:\tan\frac{SM'}{2}$$

are in a given ratio. Thus

$$\tan\frac{SM}{2}\tan\frac{SN'}{2}=\tan\frac{SM'}{2}\cdot\tan\frac{SN}{2}$$



Figure 3: Centres of similitude

is a constant. The six centres of similitude of three small circles taken in pairs lie three by three on four lines, called *axes of similitude* of the circles. Consequently if a variable circle touch two fixed circles, the line passing through the points of contact passes through a fixed point, namely, a centre of similitude of the two circles; for the points of contact are centres of similitude. Moreover if a variable circle touch two fixed circles, the tangent drawn to it from centre of similitude, through which the chord of contact passes, is constant. Thus if being given a fixed point $S$ and any curve whatever $\gamma$, on the sphere, if upon the segment of a line joining $S$ to any point $M$ of $\gamma$ a point $N'$ be taken, such that $\tan\frac{SM}{2}\tan\frac{SM'}{2}$ is constant, the locus of $N'$ is called the inverse of $\gamma$.

**Revert to the hyperbolic case** we have a new situation, namely two lines does not intersects to each other in every case. For example, if we consider three points $A, B, C$ on a line (with this order) then the ratio (by $\coth|BC| > 1$) is equal

$$\frac{\sinh|AC|}{\sinh|BC|} = \frac{\sinh(|AB|+|BC|)}{\sinh|BC|} = \cosh|AB| + \coth|BC|\sinh|AB|,$$

which is greater then $e^{|AB|}$. Therefore a ratio can be attached by a real point $C$ only if it is greater than $e^{|AB|}$ (this quantity depends on the distance of the points $A, B$). On the other hand every number greater or equal to 1 could be the ratio of hyperbolic sines of the radius of circles with origin $A$ and $B$, respectively. To solve this problem we follows the method of the book of Cyrill Vörös ([16]). We extract the concepts of measure of real elements. We expand the plane with two types of points, one type of the points at infinity and the other one the type of ideal points. With respect to a projective model these are the boundary and external points of a model with respect to the embedding real projective plane, respectively. Two parallel lines determine a point at infinity and two ultraparallel lines an ideal points which is the pole of their common transversal. Now the concept of the line can be expanded automatically; a line is real if it has real points (in this case it also has two points at infinity and the other points on it are ideal points being the poles of the real lines orthogonal to the mentioned one). The expanded real line is a closed compact set with finite length. We also distinguish the line at infinity containing precisely one point at infinity and the so-called ideal line which contains only ideal points, respectively. By definition of the common lengthes of the lines are $\pi k i$, where $k$ is a constant of the hyperbolic plane and $i$ is the imaginary unit, respectively. In this paper we assume that $k = 1$. The distance of a real and an ideal point is a complex number. Its **real part** is the distance of the real point to the polar of the ideal point with a sign, this sign is positive in the case when the polar line intersects the investigated segment and is negative otherwise. The **imaginary part** of the length is $\frac{\pi}{2}i$ implying that the sum of the lengthes of the two disjoint segments of this projective line is the total length $\pi i$. By definition the distance from a point at infinity is infinite. The distance of two ideal points of a real line is the negative of the distance of their polars. The distance of two points of an ideal line is the angle of their polars multiplied by $i$. Finally, every distances from a point at infinity of a line at infinity are undeterminable and we can define such manner that the trigonometric relationships could be valid. Thus let this distance be $\frac{\pi}{2}i$. Furthermore let the distance of two ideal points of a line at infinity be 0 or $\pi i$, accordingly that the point at infinity of the line does not belong or belongs to the corresponding segment, respectively. Now we are ready to the definition.

**Definition 2** *The* power *of a point $P$ with respect to a given cycle is the value*

$$\tanh \frac{1}{2} PA \cdot \tanh \frac{1}{2} PB$$

*, where the points $A$, $B$ are on the cycle, such that their lines passes through the point $P$. The* axe of power *of two cycles is the locus of points having the same powers with respect to the cycles. The* centres of similitude *of two cycles with non-overlapping interiors are the common points of their pairs of tangents touching direct or inverse, respectively. The first point is the* external centre of similitude *the second one is* the internal centre of similitude.

The usual statements on the euclidean or spherical power is valid also in the hyperbolic plane. The power of a point could be positive, negative or complex. (For example, in the case when the meeting points of the secant are real we have the following possibilities: positive if $P$ real and is exterior to the cycle; negative if $P$ is real and is interior to the cycle, infinity if $P$ is at infinity, or complex if $P$ is an ideal one.)

Hence we have the following six possibilities for the centres of similitude:

1. **The two cycles are circles.** To get the centres of similitude we have to solve an equation in $x$. Here $d$ means the distance of the centers of the

circles and $r \leq R$ denotes the respective radiuses.

$$\sinh(d \pm x) : \sinh x = \sinh R : \sinh r$$

from which we get that

$$\coth x = \frac{\sinh R \mp \cosh d \sinh r}{\sinh r \sinh d}$$

or equivalently

$$e^x = \sqrt{\frac{\sinh R \mp \cosh d \sinh r + \sinh r \sinh d}{\sinh R \mp \cosh d \sinh r - \sinh r \sinh d}} = \sqrt{\frac{\frac{\sinh R}{\sinh r} \mp e^d}{\frac{\sinh R}{\sinh r} + e^d}}.$$

Hence in every cases the internal centre of similitude is a real point and the external centre we have three cases. It is an ideal one, at the infinity or real point accordingly to the cases $\frac{\sinh R}{\sinh r} < e^d$, $\frac{\sinh R}{\sinh r} = e^d$ or $\frac{\sinh R}{\sinh r} > e^d$.

2. **One of the cycles is a circle and the other one is a paracycle.** The line joining the centers (which we call axe of symmetry) is a real one, but the respective ratio is zero or infinity. To determine the centres we have to decide the common tangents and their points of intersections, respectively. The external centre is ideal point and the internal centre is a real one.

3. **One of the cycles is a circle and the other one is a hypercycle.** The axe of symmetry is real the ratio of the sine of the radiuses is complex. The external centre is ideal the internal one is always real. Each of them can be determined as in the case of two circles.

4. **Each of them is a paracycle.** The axe of symmetry is real line and the internal centre is a real point. the external centre is ideal one.

5. **One of them paracycle and the other is a hypercycle.** The axe of symmetry is real line or line at infinity. The internal centre could be in the first case real point, point at infinity or ideal point, and in the second case it is ideal one. The external centre is always ideal.

6. **Both of them are hypercycles.** The axe of symmetry could be real line, ideal line or a line at infinity, and also for the internal centre we have this three possibilities. On the other hand the external centre is always ideal point.

Since two points determine a line all of the concepts using on the sphere usable on the hyperbolic plane, too. Thus we will use the concepts of "axe of similitude", "inverse and homothetic pair of points", "homothetic to" and "inverse of" a curve $\gamma$ with respect to a generalized fix point $S$ as in the case of the sphere. Thus being given a fixed point $S$ (which is the center of the cycle for which we would like to invert) and any curve whatever $\gamma$, on the plane, if upon the segment of a line joining $S$ to any point $M$ of $\gamma$ a point $N'$ be taken, such that

$$\tanh \frac{SM}{2} \tanh \frac{SN'}{2}$$

is constant, the locus of $M'$ is *called the inverse* of $\gamma$. We also use the name *cycle of inversion* for the locus of the points whose squared distance from $S$ is

$$\tanh \frac{SM}{2} \cdot \tanh \frac{SN'}{2}.$$

Among the projective elements of the pole and its polar either one of always real or both of them are at infinity. Thus in a concrete model-free construction the common point of two lines is well-defined, and in every situation it can be joined with an other point; for example, if both of them are ideal points they given by their polars (which are constructible real lines) and the required line is the polar of the intersection point of these two real lines. Thus the lengthes in the definition of the inverse can be constructed implying that inverse of a point constructible on the hyperbolic plane, too.

*Finally we remark that all of the concepts and results of inversion with respect to a sphere of the Euclidean space can be defined also in the hyperbolic space, the "basic sphere" could be hypersphere, parasphere or sphere, respectively. As in the plane we can use also the concept of ideal elements and the concept of elements at infinity, if it is necessary. It can be proved (using Poincare's ball-model) that every hyperbolic plane of the hyperbolic space can be inverted to a sphere by such a general inversion. This map sends the cycles of the plane to circles of the sphere.*

### 1.3.2 Steiner's construction on the hyperbolic plane

**Theorem 2** *The Steiner's construction can be done also in the hyperbolic plane. More precisely for three given non-overlapping cycles can be constructed three other, each of them touching the two others and also two of the three given.*

**Proof:** Denote by $c_i$ the given cycles. Now the steps of Steiner's construction are the followings:

1. Construct the cycle of inversion $c_{i,j}$, for the given cycles $c_i$ and $c_j$, where the center of inversion is the external centre of similitude of them.
2. Construct cycles $k_j$ touching two cycles $c_{i,j}, c_{j,k}$ and the given cycle $c_j$.
3. Construct the cycles $l_{i,j}$ touching $k_i$ and $k_j$ through the point $P_k = k_k \cap c_k$.
4. Construct the Malfatti's cycles $m_j$ as the common touching cycle of the four cycles $l_{i,j}$, $l_{j,k}$, $c_i$, $c_k$.

From the definitions of the preceding paragraph we know that the first step is constructible.

To prove the constructible of the second step we interpret Gergonne's construction (see e.g. [8]) for hyperbolic plane. This construction is:

*Draw the point of power of the given circles and an axe of similitude of three centres of similitude, respectively. The poles of the axe of similitude with respect to the given circles are points of the rays originated from the point of power of the three circle and passes through the touching points of a touching circle and one of the three given ones, respectively.*

We note that by this method we find the common touching circles pairwise, correspondingly to the choose of the axe of similitude. In our problem we have to take into consideration only one axe of similitude and the corresponding touching cycles. As we saw, all of the using concept works on the hyperbolic plane, too. If we consider a pair of resulting cycles touching the three given ones then the line joining the touching points passes through one of their centre of similitude. Thus this centre of similitude of the sought cycles is the point of power of the given triplet. On the other hand two of the three given cycles are a touching pair with respect to the sought cycles, hence its centre of similitude has the same power with respect to the sought cycles. So the three centres of similitude are on the axe of power of the sought cycles hence it is also the axe of similitude with respect to the given cycles, say $s$. Since the pole (with respect to a given cycle) of the line joining the corresponding two touching points is also on $s$, by the theorem of pole-polar we get that the pole of $s$ lies on the mentioned secant. This proves the construction.

The third step in construction is a trivialization of the second step. (A given cycle is a point now.) Obviously the general construction can be done also in this case.

The fourth step is again the second one choosing three arbitrary cycles from the four ones if the quadrangles determined the cycles are inscriptible.

Finally we have to prove that this construction gives the Malfatti's cycles. As we saw the Malfatti's cycles are exist (see Theorem 1). We also know that in an embedding hyperbolic space the examined plane can be inverted to a sphere. The trigonometry of the sphere is absolute implying that the possibility of a construction which can be checked by trigonometric calculations, independent to the fact that the embedding space was hyperbolic or Euclidean one. Of course, the Steiner's construction just such a like, to touching of circles of the sphere can be checked by spherical trigonometry. So we may assume that the examined sphere is a sphere of the Euclidean space and we can apply Cayley's analytical research (see in [3]) in which he proved that Steiner's construction works for a surface of second order. Hence the above construction produces the required touches. □

## 2 A note on Malfatti's marble problem

As we can read in the paper [1] there is a long history of Malfatti's marble problem which is: *"Given a triangle find three non-overlapping circles inside it of total maximum area."* Also in this paper we can find an elegant proof for the theorem: *"The greedy arrangement has the largest total area among of non-overlapping circles in a triangle."*

The apropos that I join the marble problem the fact that the hint for the proof of Remark 2 in [1] is insufficient. As we can see from the foregoing proof the mentioned absolute lemma on quadrilateral is only one point needs to be checked since the value $\frac{r_1+r_2}{2}$ is not the radius of the touching circle at the midpoint $F_1$ of the segment $A_1A_2$ (see in Fig. 4).

**Theorem 3** *On the hyperbolic plane the greedy arrangement has the largest total area among of non-overlapping circles in a triangle.*



Figure 4: Comparison of two circles rigid arrangements on the hyperbolic plane

**Proof:**

$$|F_1F_2| \leq \frac{|A_1B_1| + |A_2B_2|}{2} = \frac{(r_1+r_2) + (R(r_1)+R(r_2))}{2}$$

On the other hand on the hyperbolic plain we have

$$|F_1T_1| \leq \frac{r_1+r_2}{2} \text{ and } |F_2T_2| \leq \frac{R(r_1)+R(r_2)}{2}.$$

Let now $F_i'$ such that the distance segments $F_i'T_i'$ hold the equalities

$$|F_1'T_1'| = \frac{r_1+r_2}{2} \text{ and } |F_2'T_2'| = \frac{R(r_1)+R(r_2)}{2},$$

respectively. If the bisectors $O_1A_1$, $O_2A_2$ intersect each other in $M$, then we have

$$(O_1A_iM), (O_2B_iM) \text{ for } i = 1, 2$$

which are the respective orders of points on their lines. This implies that the orders

$$(F_1F_1'M), (F_2F_2'M)$$

are also valid, hence

$$|F_1'F_2'| \leq |F_1F_2| \leq \frac{r_1+r_2+R(r_1)+R(r_2)}{2} = |F_1'T_1'| + |F_2'T_2'|$$

is also true. This implies that

$$R\left(\frac{r_1+r_2}{2}\right) = R(F_1'T_1') \leq |F_2'T_2'| = \frac{R(r_1)+R(r_2)}{2}$$

showing that $R$ is a convex function. Since the area of a hyperbolic circle as the function of its radius is $2\pi(\cosh r - 1)$ an increasing convex function, too the Bezdek's and at all proof valid in also the hyperbolic plane as they stated in [1]. □

We note that this proof is valid in the case of asymptotic triangle's, too. The statements on concave triangles are correct. We note that the corresponding proof for the case when among the vertices have points at infinity works, too.

# References

[1] Andreatta M., Bezdek A., Boronski J.P., *The problem of Malfatti: Two Centuries of Debate.* (to appear in the Mathematical Intelligencer) (published online 13 July 2010).

[2] Bottema O., *The Malfatti Problem.* Forum Geometricorum **1** (2001) 43–50.

[3] Cayley, A.,*Analytical Researches Connected with Steiner's Extension of Malfatti's Problem* Phil. Trans. of the Roy. Soc. of London, **142** (1852), 253–278.

[4] Cayley, A.,*On Schellbachs solution of Malfattis problem* Quart. J. Pure Appl. Math. **1** (1857) 222226; CMP III (1890) 4447.

[5] Cayley, A.,*On a System of Equations connected with Malfatti's Problem and on another Algebraic System* Camb. and Dubl. Math. Journ. **4** (1849) 270 [79].

[6] Casey, J., *A sequel to the First Six Books of the Elements of Euclid, Containing an Easy Introduction to Modern Geometry with Numerous Examples, 5th. ed.,* Hodges, Figgis and Co., Dublin (1888).

[7] Coxeter, H.S.M., Non Euclidean Geometry Toronto, 1947.

[8] Dörrie, H., Triumph der mathematik Physica-Verlag, Würzburg, 1958.

[9] Hart, A. S., *Geometric investigations of Steiner's construction for Malfatti's problem.* Quart. J. Pure Appl. Math. **1** (1857) 219221.

[10] Malfatti, G., *Memoria sopra un problema sterotomico* Memorie dimatematica e di Fisica della Societa Italiano delle Scienze, **10** (1803), 235-244.

[11] Molnár, E., *Inversion auf der idealebene der Bachmannschen metrischen ebene* Acta Math. Acad. Sci. Hungar. **37/4** (1981), 451–470.

[12] Schellbach, *Eine Lösung der Malfattischen Aufgabe* Journal fr die reine und angewandte Mathematik **45/1** (1853) 91-92.

[13] Schellbach, *Eine Erweiterung der Malfattischen Aufgabe* Journal fr die reine und angewandte Mathematik **45/2** (1853) 186-187.

[14] Steiner's gesammelte werke, (Herausgegeben K. Weierstrass) Berlin, 1881.

[15] Steiner, J., *Einige geometrische Betrachtungen.* Journal fr die reine und angewandte Mathematik **1/2** (1826) 161-184, **1/3** (1826) 252-288.

[16] Dr. Vörös, C., Analytic Bolyai's geometry, Budapest, 1909 (in hungarian).

Ákos G.Horváth
Department of Geometry, Mathematical Institute
Budapest University of Technology and Economics,
H-1521 Budapest,
Hungary
e-mail: ghorvath@math.bme.hu